

Tilburg University

A characterization of the value of zero-sum two person games

Tijs, S.H.

Published in:
Naval Research Logistics

Publication date:
1981

[Link to publication in Tilburg University Research Portal](#)

Citation for published version (APA):
Tijs, S. H. (1981). A characterization of the value of zero-sum two person games. *Naval Research Logistics*, 28, 153-156.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

A CHARACTERIZATION OF THE VALUE OF ZERO-SUM TWO-PERSON GAMES

S. H. Tijs

*Department of Mathematics
Catholic University
Nijmegen, The Netherlands*

ABSTRACT

For the family D , consisting of those zero-sum two-person games which have a value, the value-function on D is characterized by four properties called objectivity, monotony, symmetry and sufficiency.

INTRODUCTION

In a beautiful paper, E. I. Vilkas gave a characterization of the value-function, defined on the class of all finite matrix games [2]. In [1], pp. 60-65 this result was extended to the class of all finite and semi-infinite matrix games.

The purpose of this paper is to deduce characterizing properties for the value-function on the set of all determined two-person games. The organization of the paper is as follows: the necessary notation and definitions are given in sections 1 and 2; in section 3, properties for the value-function are presented, which are shown in section 4 to be characteristic of this function.

1. A (zero-sum) two-person game is an ordered triple $\langle X, Y, K \rangle$, in which X and Y are nonempty sets (called the *pure strategy spaces* of player I and player II, respectively) and $K : X \times Y \rightarrow \mathbb{R}$ is a real-valued function on the Cartesian product of X and Y (called the *pay off function* of player I).

2. Let $\langle X, Y, K \rangle$ be a two-person game. For each $x \in X$ ($y \in Y$) let us denote the probability measure on X (Y) with mass 1 in x (y) by e_x (e_y). Let P_X be the set of all convex combinations of elements of $\{e_x; x \in X\}$; likewise let P_Y be the convex hull of $\{e_y; y \in Y\}$. Then the two-person game $\langle P_X, P_Y, E_K \rangle$ with

$$E_K(\mu, \nu) := \int \int K(x, y) d\mu(x) d\nu(y) \text{ for each } (\mu, \nu) \in P_X \times P_Y$$

is called the *c-mixed extension* of the game $\langle X, Y, K \rangle$. The lower value $\sup_{\mu \in P_X} \inf_{\nu \in P_Y} E_K(\mu, \nu)$ of the game $\langle P_X, P_Y, E_K \rangle$ is denoted by $\underline{v}(X, Y, K)$ and the upper value $\inf_{\nu \in P_Y} \sup_{\mu \in P_X} E_K(\mu, \nu)$ is denoted by $\bar{v}(X, Y, K)$. Note that

$$-\infty \leq \underline{v}(X, Y, K) \leq \bar{v}(X, Y, K) \leq \infty.$$

If $\underline{v}(X, Y, K) = \bar{v}(X, Y, K)$ for a game $\langle X, Y, K \rangle$, then we say that the game is a *determined game*. In that case, the common value is denoted by $v(X, Y, K)$ and called the *value* of (the c-mixed extension of) $\langle X, Y, K \rangle$. The family of determined games is denoted by D .

3. In this section we want to look at some distinguished properties of the value-function $v : D \rightarrow [-\infty, \infty]$. For this purpose we need some definitions.

DEFINITION 1: The *transpose* of a two-person game $\langle X, Y, K \rangle$ is the two-person game $\langle Y, X, -K' \rangle$ where

$$K'(y, x) := K(x, y) \text{ for each } (y, x) \in Y \times X.$$

DEFINITION 2: Let $\langle X, Y, K \rangle$ be a two-person game and let S be a nonempty subset of X . Then we say that S is *sufficient for player I in the game* $\langle X, Y, K \rangle$ if for each $x \in X - S$ there exists a $\mu \in P_S$ such that

$$E_K(\mu, e_y) \geq K(x, y) \text{ for each } y \in Y.$$

DEFINITION 3: Let $\langle X, Y, K \rangle$ be a two-person game and let T be a nonempty subset of Y . We say that T is *sufficient for player II in the game* $\langle X, Y, K \rangle$ if T is sufficient for player I in the game $\langle Y, X, -K' \rangle$.

THEOREM 1:

(P.1) ["Objectivity"] Let $\langle X, Y, K \rangle$ be a two-person game and suppose that $X = \{a\}$, $Y = \{b\}$. Then $\langle X, Y, K \rangle \in D$ and $v(X, Y, K) = K(a, b)$.

(P.2) ["Monotonicity"] Let $\langle X, Y, K \rangle \in D$ and $\langle X, Y, L \rangle \in D$ and suppose that $L \geq K$ (i.e. $L(x, y) \geq K(x, y)$ for each $(x, y) \in X \times Y$). Then $v(X, Y, L) \geq v(X, Y, K)$.

(P.3) ["Symmetry"] Let $\langle X, Y, K \rangle \in D$. Then $\langle Y, X, -K' \rangle \in D$ and $v(Y, X, -K') = -v(X, Y, K)$.

(P.4) ["Sufficiency"] Let $\langle X, Y, K \rangle$ be a two-person game, and $\emptyset \neq S \subset X$ and let $K' : S \times Y \rightarrow \mathbb{R}$ be the restriction of K to $S \times Y$. Suppose that S is sufficient for player I in the game $\langle X, Y, K \rangle$. Then $\langle S, Y, K' \rangle \in D$ iff $\langle X, Y, K \rangle \in D$ and

$$v(X, Y, K) = v(S, Y, K') \text{ if } \langle S, Y, K' \rangle \in D.$$

PROOF: (P.1) and (P.2) are obvious. (P.3) follows from the fact that

$$-E_K(\mu, \nu) = E_{-K'}(\nu, \mu) \text{ for each } (\mu, \nu) \in P_X \times P_Y.$$

Now let us prove (P.4). First we note that P_S can be seen (in an obvious manner) as a subset of P_X , and that $E_{K'}$ is the restriction of E_K to $P_S \times P_Y$.

Take $\alpha \in P_X$. Then there exist $n \in \mathbb{N}$, $x^1, x^2, \dots, x^n \in X$ and $p_1, p_2, \dots, p_n \in [0, \infty)$ such that $\sum_{i=1}^n p_i = 1$ and $\alpha = \sum_{i=1}^n p_i e_{x^i}$. Since S is sufficient for player I in the game $\langle X, Y, K \rangle$ for each $i \in \{1, \dots, n\}$, there exists an $\alpha_i \in P_S$ such that

$$E_K(\alpha_i, e_y) \geq K(x^i, y) \text{ for each } y \in Y.$$

[If $x^i \in S$, then we can take $\alpha_i = e_{x^i}$.] Let $\bar{\alpha} := \sum_{i=1}^n p_i \alpha_i$. Then $\bar{\alpha} \in P_S$ and

$$E_K(\bar{\alpha}, e_y) = \sum_{i=1}^n p_i E_K(\alpha_i, e_y) \geq \sum_{i=1}^n p_i K(x^i, y) = E_K(\alpha, e_y) \text{ for each } y \in Y.$$

But then,

$$(i) \quad E_K(\bar{\alpha}, \nu) \geq E_K(\alpha, \nu) \text{ for each } \alpha \in P_X \text{ and each } \nu \in P_Y.$$

This implies that

$$\sup_{\mu \in P_S} E_{K'}(\mu, \nu) = \sup_{\alpha \in P_X} E_K(\alpha, \nu) \text{ for each } \nu \in P_Y$$

and thus

$$(ii) \quad \bar{v}(S, Y, K') = \bar{v}(X, Y, K).$$

From (i) we may also conclude that

$$\inf_{\nu \in P_Y} E_{K'}(\bar{\alpha}, \nu) \geq \inf_{\nu \in P_Y} E_K(\alpha, \nu) \text{ for each } \alpha \in P_X$$

and then

$$(iii) \quad \underline{v}(S, Y, K') = \underline{v}(X, Y, K).$$

Now (P.4) follows from (ii) and (iii). ||

4. The following theorem shows that the properties (P.1)-(P.4) characterize the value-function $v : D \rightarrow [-\infty, \infty]$.

THEOREM 2: Let $f : D \rightarrow [-\infty, \infty]$ be a function with the following four properties:

(Q.1) If $X = \{a\}$, $Y = \{b\}$ and if K is a real-valued function on $X \times Y$, then $f(X, Y, K) = K(a, b)$.

(Q.2) For each $\langle X, Y, K \rangle \in D$, $\langle X, Y, L \rangle \in D$ with $L \geq K$: $f(X, Y, L) \geq f(X, Y, K)$.

(Q.3) For each $\langle X, Y, K \rangle \in D$: $f(Y, X, -K') = -f(X, Y, K)$.

(Q.4) For each $\langle X, Y, K \rangle \in D$ and $\langle S, Y, K' \rangle \in D$, where $S \subset X$, K' is the restriction of K to $S \times Y$ and where S is sufficient for player I in the game $\langle X, Y, K \rangle$, we have $f(S, Y, K') = f(X, Y, K)$.

Then $f(X, Y, K) = v(X, Y, K)$ for each $\langle X, Y, K \rangle \in D$.

PROOF: First we note that (Q.3) and (Q.4) imply

(Q.5) For each $\langle X, Y, K \rangle \in D$ and $\langle X, T, K'' \rangle \in D$, where $T \subset Y$, K'' is the restriction of K to $X \times T$ and where T is sufficient for player II in the game $\langle X, Y, K \rangle$, we have $f(X, T, K'') = f(X, Y, K)$.

Now take an $\langle X, Y, K \rangle \in D$ with $v(X, Y, K) \in (-\infty, \infty]$ and take a real number t such that $v(X, Y, K) > t$. We want to prove that $f(X, Y, K) \geq t$. For this purpose we introduce the following five two-person games:

(1) $\langle X \cup \{a\}, Y, L \rangle$ where $a \notin X$ and where $L(x, y) := K(x, y)$ for each $(x, y) \in X \times Y$ and $L(a, y) := t$ for each $y \in Y$.

(2) $\langle X \cup \{a\}, Y, M \rangle$ where $M(x, y) := \text{minimum } \{K(x, y), t\}$ for each $(x, y) \in (X \cup \{a\}) \times Y$.

- (3) $\langle X \cup \{a\}, Y \cup \{b\}, N \rangle$ where $b \notin Y$ and where $N(x, y) := M(x, y)$ for each $(x, y) \in (X \cup \{a\}) \times Y$ and $N(x, b) := t$ for each $x \in X \cup \{a\}$.
- (4) $\langle \{a\}, Y \cup \{b\}, N' \rangle$ where N' is the restriction of N to $\{a\} \times (Y \cup \{b\})$.
- (5) $\langle \{a\}, \{b\}, N'' \rangle$ where N'' is the restriction of N' to $\{a\} \times \{b\}$.

Since $v(X, Y, K) > t$, there exists a $\mu \in P_X$ such that

$$E_L(\mu, e_y) = E_K(\mu, e_y) \geq t = L(a, y) \text{ for each } y \in Y.$$

Hence, X is sufficient for player I in the game $\langle X \cup \{a\}, Y, L \rangle$. By (P.4) and (Q.4) we may conclude that

$$(Q.6) \quad \langle X \cup \{a\}, Y, L \rangle \in D \text{ and } f(X \cup \{a\}, Y, L) = f(X, Y, K).$$

It follows from (Q.1) that

$$(Q.7) \quad f(\{a\}, \{b\}, N'') = t.$$

In the game $\langle \{a\}, Y \cup \{b\}, N' \rangle$ the set $\{b\}$ is sufficient for player II because

$$E_{N'}(e_a, e_b) = N'(a, y) = t \text{ for each } y \in Y.$$

Then $\langle \{a\}, Y \cup \{b\}, N' \rangle \in D$ in view of (P.3) and (P.4); now (Q.5) implies

$$(Q.8) \quad f(\{a\}, Y \cup \{b\}, N') = f(\{a\}, \{b\}, N'').$$

In the game $\langle X \cup \{a\}, Y \cup \{b\}, N \rangle$ the set $\{a\}$ is sufficient for player I because for each $x \in X$:

$$t = E_N(e_a, e_y) \geq N(x, y) \text{ for each } y \in Y \cup \{b\}.$$

By (P.4) and (Q.4) we obtain: $\langle X \cup \{a\}, Y \cup \{b\}, N \rangle \in D$ and

$$(Q.9) \quad f(X \cup \{a\}, Y \cup \{b\}, N) = f(\{a\}, Y \cup \{b\}, N').$$

It is easy to see that Y is sufficient for player II in the game $\langle X \cup \{a\}, Y \cup \{b\}, N \rangle$. Hence, by (P.3) and (P.4): $\langle X \cup \{a\}, Y, M \rangle \in D$; and then by (Q.5):

$$(Q.10) \quad f(X \cup \{a\}, Y, M) = f(X \cup \{a\}, Y \cup \{b\}, N).$$

Now $L \geq M$ and then by (Q.2) we have

$$(Q.11) \quad f(X \cup \{a\}, Y, L) \geq f(X \cup \{a\}, Y, M).$$

Combining (Q.6)-(Q.11) we obtain: $f(X, Y, K) \geq t$. Thus, we have proved that $f(X, Y, K) \geq t$ for each $\langle X, Y, K \rangle \in D$ with $v(X, Y, K) \in (-\infty, \infty]$ and each $t < v(X, Y, K)$. But then

$$(Q.12) \quad f(X, Y, K) \geq v(X, Y, K) \text{ for each } \langle X, Y, K \rangle \in D.$$

It follows from (Q.3), (Q.12) and (P.3) that

$$(Q.13) \quad f(X, Y, K) = -f(Y, X, K') \leq -v(Y, X, K') = v(X, Y, K) \text{ for each } \langle X, Y, K \rangle \in D.$$

Properties (Q.12) and (Q.13) imply the conclusion of the theorem. \parallel

REFERENCES

- [1] Tijds, S.H., *Semi-Infinite and Infinite Matrix Games and Bimatrix Games*, Ph.D. dissertation, Department of Mathematics, Catholic University, Nijmegen, The Netherlands (1975).
- [2] Vilkas, E.I., "Axiomatic Definition of the Value of a Matrix Game," *Theory of Probability and its Applications* 8, 304-307 (1963).